

MEDICAL POLICY

POLICY TITLE	INJECTABLE BULKING AGENTS FOR THE TREATMENT OF URINARY AND FECAL INCONTINENCE
POLICY NUMBER	MP 4.012

CLINICAL BENEFIT	<input checked="" type="checkbox"/> MINIMIZE SAFETY RISK OR CONCERN. <input checked="" type="checkbox"/> MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. <input type="checkbox"/> ASSURE APPROPRIATE LEVEL OF CARE. <input type="checkbox"/> ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. <input type="checkbox"/> ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. <input type="checkbox"/> ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
Effective Date:	4/1/2024

[POLICY RATIONALE](#)
[DISCLAIMER](#)
[POLICY HISTORY](#)

[PRODUCT VARIATIONS](#)
[DEFINITIONS](#)
[CODING INFORMATION](#)

[DESCRIPTION/BACKGROUND](#)
[BENEFIT VARIATIONS](#)
[REFERENCES](#)

I. POLICY

The use of carbon-coated spheres (Durasphere®), calcium hydroxylapatite (CaHA) (Coaptite®), polyacrylamide hydrogel (Bulkamid®), or polydimethylsiloxane (Macroplastique®) may be considered **medically necessary** to treat urinary stress incontinence in men and women who have failed conservative therapy.

The use of autologous cellular therapy (e.g., myoblasts, fibroblasts, muscle-derived stem cells, or adipose-derived stem cells), autologous fat, and autologous ear chondrocytes to treat stress urinary incontinence is considered **investigational**.

The use of any other periurethral bulking agent, including, but not limited to Teflon®, to treat stress urinary incontinence is considered **investigational**.

The use of periurethral bulking agents to treat urge urinary incontinence is considered **investigational**.

The use of perianal bulking agents to treat fecal incontinence is considered **investigational**. There is insufficient evidence to support a general conclusion concerning the health outcomes related to their efficacy for the above listed investigational indications.

Policy Guidelines

Individuals should have had inadequate response to conservative therapy or therapies; in general, these treatments should have been used for at least 3 months. Conservative therapy for stress incontinence includes pelvic floor muscle exercises and behavioral changes, such as fluid management and moderation of physical activities that provoke incontinence. Additional options include intravaginal estrogen therapy, use of a pessary, and treatment of other underlying causes of incontinence in patients amenable to these treatments.

Cross-references

- MP 1.033** Sacral Nerve Modulation/Stimulation and Pelvic Floor Stimulation Devices
- MP 1.134** Percutaneous Tibial Nerve Stimulation

MEDICAL POLICY

POLICY TITLE	INJECTABLE BULKING AGENTS FOR THE TREATMENT OF URINARY AND FECAL INCONTINENCE
POLICY NUMBER	MP 4.012

MP 2.064 Biofeedback and Neurofeedback Therapy

MP 4.034 Transvaginal and Transurethral Radiofrequency Tissue Remodeling for Urinary Stress Incontinence

II. PRODUCT VARIATIONS

[Top](#)

This policy is only applicable to certain programs and products administered by Capital Blue Cross please see additional information below, and subject to benefit variations as discussed in Section VI below.

FEP PPO - Refer to FEP Medical Policy Manual. The FEP Medical Policy Manual can be found at: <https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>.

III. DESCRIPTION/BACKGROUND

[Top](#)

Incontinence

Incontinence, especially urinary, is a common condition and can have a substantial impact on quality of life. Estimates from the National Center for Health Statistics have suggested that, among noninstitutionalized persons 65 years of age and older, 44% have reported issues with urinary incontinence and 17% issues with fecal incontinence.

Treatment

Urinary Incontinence

Injectable bulking agents are space-filling substances used to increase tissue bulk. When used to treat stress urinary incontinence (SUI), bulking agents are injected periurethrally to increase tissue bulk and thereby increase resistance to the outflow of urine. The bulking agent is injected into the periurethral tissue as a liquid that then solidifies into a spongy material to bulk the urethral wall. Bulking agents may be injected over a course of several treatments until the desired effect is achieved. Periurethral bulking agents have been widely used for incontinence in women. Men have also been treated, typically those with post-prostatectomy incontinence.

Key factors in determining the optimal product are biocompatibility, durability, and absence of migration. A number of periurethral bulking agents to treat urinary incontinence have been cleared for marketing by the Food and Drug Administration (FDA); however, products developed to date have not necessarily met all criteria of the ideal bulking agents. The first FDA-approved product was cross-linked collagen (e.g., Contigen). The agent was found to be absorbed over time and symptoms could recur, requiring additional injections. Contigen production was discontinued in 2011. Other periurethral bulking agents cleared by FDA for urinary incontinence include carbon-coated beads (e.g., Durasphere), spherical particles of calcium hydroxylapatite (CaHA®) in a gel carrier (Coaptite®), polydimethylsiloxane (silicone, Macroplastique®), cross-linked polyacrylamide hydrogel (Bulkamid®), and ethylene vinyl alcohol copolymer implants (e.g., Tegress®, formerly Uryx®). Tegress was voluntarily removed from the market due to safety concerns.

MEDICAL POLICY

POLICY TITLE	INJECTABLE BULKING AGENTS FOR THE TREATMENT OF URINARY AND FECAL INCONTINENCE
POLICY NUMBER	MP 4.012

Fecal Incontinence

After the success of periurethral bulking agents for treating SUI, bulking agents injected into the anal canal have been proposed for treating fecal incontinence. In particular, bulking agents are a potential treatment for passive fecal incontinence associated with internal anal sphincter (IAS) dysfunction. The bulking agent is injected into the submucosa of the anal canal to increase tissue bulk in the area, which narrows the opening of the anus. Current treatment options for fecal incontinence include conservative measures e.g., dietary changes, pharmacotherapy and pelvic floor muscle exercises, sacral nerve stimulation, and surgical interventions to correct an underlying problem.

Several agents identical or similar to those used for urinary incontinence (e.g., Durasphere, silicone biomaterial) have been studied for the treatment of fecal incontinence. To date, only one bulking agent has been approved by the FDA for treating fecal incontinence. This formulation of non-animal stabilized hyaluronic acid/dextranomer in stabilized hyaluronic acid (NASHA Dx) and is marketed by Q-Med as Solesta. A hyaluronic acid/dextranomer formulation (Deflux™) from the same company has been commercially available for a number of years for the treatment of vesicoureteral reflux in children.

Autologous fat and autologous ear chondrocytes have also been used as periurethral bulking agents; autologous substances do not require FDA approval. Polytetrafluoroethylene (Teflon®) has been investigated as an implant material but has not received FDA approval. A more recently explored alternative is cellular therapy with myoblasts, fibroblasts, or stem cells (muscle-derived or adipose-derived). In addition to their use as periurethral bulking agents, it is hoped that transplanted stem cells will undergo self-renewal and multipotent differentiation, which could result in regeneration of the sphincter and its neural connections.

In May, 2023, the American Society of Colon and Rectal Surgeons published Clinical Practice Guidelines for the Management of Fecal Incontinence. The guideline states that “injection of biocompatible bulking agents into the anal canal is not routinely recommended for the treatment of FI (fecal incontinence)” citing limited improvement over placebo, diminishing long-term results, and cost.

Regulatory Status

Several periurethral bulking agents have been approved by the FDA through the premarket approval process for the treatment of SUI due to intrinsic sphincter deficiency; other than Contigen®, approval is only for use in adult women. Products include:

- In 1993, Contigen (Allergan, Inc.), a cross-linked collagen, was approved. A supplemental approval in 2009 limited the device’s indication to treatment of urinary incontinence due to intrinsic sphincter deficiency in patients (men or women) who have shown no improvement in incontinence for at least 12 months. Allergan ceased production in 2011; no reason for discontinuation was provided to the public.
- In 1999, Durasphere (Advanced UroScience), pyrolytic carbon-coated zirconium oxide spheres, was approved.
- In 2004, Uryx (CR Bard), vinyl alcohol copolymer implants, was approved. In 2005, approval was given to market the device under the trade name Tegress. In 2007, Tegress was voluntarily removed from the market due to safety concerns.

MEDICAL POLICY

POLICY TITLE	INJECTABLE BULKING AGENTS FOR THE TREATMENT OF URINARY AND FECAL INCONTINENCE
POLICY NUMBER	MP 4.012

- In 2005, Coaptite (Merz Aesthetics, previously BioForm Medical), spherical particles of calcium hydroxylapatite, suspended in a gel carrier, was approved
- In 2006, Macroplastique (Cogentix Medical), polydimethylsiloxane, was approved.
- In 2020, Bulkamid Urethral Bulking System (Axonics Modulation Technologies, Inc.), a soft hydrogel that consists of 97.5% water and 2.5% polyacrylamide, was approved.

In 2011, NASHA Dx, marketed as Solesta® (Q-Med), was approved by FDA through the premarket approval process as a bulking agent to treat fecal incontinence in patients 18 years and older who have failed conservative therapy. FDA product code: LNM.

IV. RATIONALE

[Top](#)

Summary of Evidence

For individuals who have SUI who receive injectable bulking agents, the evidence includes randomized controlled trials (RCTs) and systematic reviews of RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The trials vary by bulking agents used and comparator interventions (eg, placebo, conservative therapy, surgical procedure, another bulking agent). Due to this heterogeneity across studies, and the small number of studies in each category, Cochrane reviewers were unable to draw specific conclusions about the efficacy of specific bulking agents compared with alternative treatments. Additionally, authors of another recent systematic review concluded that bulking agents were less effective than surgical procedures regarding subjective improvement after treatment, with no difference between the interventions with regard to complications. Studies have shown that cross-linked collagen improves the net health outcome (i.e., it is effective in some patients who failed conservative treatment with fewer adverse events than surgery), although products that cross-link in such a way are no longer commercially available. There is evidence that Food and Drug Administration (FDA) FDA-approved carbon-coated spheres, calcium hydroxylapatite, polyacrylamide hydrogel and, and polydimethylsiloxane have efficacy for treating incontinence and produce outcomes and have a safety profile similar to cross-linked collagen. The evidence is sufficient to determine qualitatively that the technology results in a meaningful improvement in the net health outcome.

For individuals who have fecal incontinence who receive injectable bulking agents, the evidence includes RCTs and systematic reviews of RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. A comparative effectiveness review from the Agency for Healthcare Research and Quality evaluated 2 RCTs with the FDA-approved product NASHA Dx (Solesta) and 2 RCTs with Durasphere (off-label in the United States). One RCT comparing NASHA Dx with sham found that NASHA Dx improved some outcome measures but not others. The other RCT did not find a significant difference in efficacy between NASHA Dx and biofeedback. Two additional RCTs evaluating Durasphere found only short-term improvements in fecal incontinence severity. Controlled trials with longer follow-up are important to determine the durability of any treatment effect. The evidence is insufficient to determine the effects of the technology on health outcomes.

MEDICAL POLICY

POLICY TITLE	INJECTABLE BULKING AGENTS FOR THE TREATMENT OF URINARY AND FECAL INCONTINENCE
POLICY NUMBER	MP 4.012

V. DEFINITIONS

[Top](#)

510(k) APPROVAL refers to section 510(k) of the Food, Drug and Cosmetic Act. Under 510(k), before a manufacturer can market a medical device in the United States, they must demonstrate to FDA's satisfaction that it is substantially equivalent (as safe and effective) to a device already on the market.

MIXED INCONTINENCE is a combination of stress and urge incontinence.

OVERFLOW INCONTINENCE is characterized by small frequent voiding due to overfilling of the bladder or to a bladder with pathologically decreased volume.

PESSARY is a device inserted into the vagina to function as a support structure for the uterus.

STRESS INCONTINENCE is an involuntary loss of urine that occurs during physical activity, such as coughing, sneezing, laughing or exercise. This incontinence occurs as a result of weakened pelvic muscles that support the bladder and urethra, or because of malfunction of the urethral sphincter.

URGE INCONTINENCE is a condition characterized by a strong desire to urinate immediately before an involuntary bladder contraction with a loss of a large amount of urine.

VI. BENEFIT VARIATIONS

[Top](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's health benefit plan. Benefit determinations should be based in all cases on the applicable health benefit plan language. Medical policies do not constitute a description of benefits. A member's health benefit plan governs which services are covered, which are excluded, which are subject to benefit limits and which require preauthorization. There are different benefit plan designs in each product administered by Capital Blue Cross. Members and providers should consult the member's health benefit plan for information or contact Capital Blue Cross for benefit information.

VII. DISCLAIMER

[Top](#)

Capital Blue Cross's medical policies are developed to assist in administering a member's benefits, do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member's benefit information, the benefit information will govern. If a provider or a member has a question concerning the application of this medical policy to a specific member's plan of benefits, please contact Capital Blue Cross' Provider Services or Member Services. Capital Blue Cross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.

VIII. CODING INFORMATION

[Top](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is

MEDICAL POLICY

POLICY TITLE	INJECTABLE BULKING AGENTS FOR THE TREATMENT OF URINARY AND FECAL INCONTINENCE
POLICY NUMBER	MP 4.012

determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Investigational; therefore, not covered:

Procedure Codes								
L8605								

Covered when medically necessary:

Procedure Codes								
L8606	51715							

ICD-10-CM Diagnosis Code	Description
N36.42	Intrinsic sphincter deficiency (ISD)
N36.43	Combined hypermobility of urethra and intrinsic sphincter deficiency
N36.8	Other specified disorders of urethra
N39.3	Stress incontinence (female) (male)
N39.4	Other specified urinary incontinence
N39.46	Mixed incontinence (urge and stress incontinence)

IX. REFERENCES

[Top](#)

1. Gorina Y, Schappert S, Bercovitz A, et al. Prevalence of incontinence among older americans. *Vital Health Stat 3*. Jun 2014; (36): 1-33. PMID 24964267
2. Agency for Health Care Policy and Research. *Clinical Practice Guideline. Urinary Incontinence in Adults*. Rockville, MD: Department of Health and Human Services; 1996.
3. Corcos J, Collet JP, Shapiro S, et al. Multicenter randomized clinical trial comparing surgery and collagen injections for treatment of female stress urinary incontinence. *Urology*. May 2005; 65(5): 898-904. PMID 15882720
4. Hoe V, Haller B, Yao HH, et al. Urethral bulking agents for the treatment of stress urinary incontinence in women: A systematic review. *Neurourol Urodyn*. Aug 2021; 40(6): 1349-1388. PMID 34015151
5. Pivazyan L, Kasyan G, Grigoryan B, et al. Effectiveness and safety of bulking agents versus surgical methods in women with stress urinary incontinence: a systematic review and meta-analysis. *Int Urogynecol J*. Aug 05 2021. PMID 34351463
6. Kirchin V, Page T, Keegan PE, et al. Urethral injection therapy for urinary incontinence in women. *Cochrane Database Syst Rev*. Jul 25 2017; 7: CD003881. PMID 28738443
7. Kirchin V, Page T, Keegan PE, et al. Urethral injection therapy for urinary incontinence in women. *Cochrane Database Syst Rev*. Feb 15 2012; (2): CD003881. PMID 22336797

MEDICAL POLICY

POLICY TITLE	INJECTABLE BULKING AGENTS FOR THE TREATMENT OF URINARY AND FECAL INCONTINENCE
POLICY NUMBER	MP 4.012

8. Davila GW. Nonsurgical outpatient therapies for the management of female stress urinary incontinence: long-term effectiveness and durability. *Adv Urol.* 2011; 2011: 176498. PMID 21738529
9. Lightner D, Calvosa C, Andersen R, et al. A new injectable bulking agent for treatment of stress urinary incontinence: results of a multicenter, randomized, controlled, double-blind study of Durasphere. *Urology.* Jul 2001; 58(1): 12-5. PMID 11445471
10. Mayer RD, Dmochowski RR, Appell RA, et al. Multicenter prospective randomized 52-week trial of calcium hydroxylapatite versus bovine dermal collagen for treatment of stress urinary incontinence. *Urology.* May 2007; 69(5): 876-80. PMID 17482925
11. Sokol ER, Karram MM, Dmochowski R. Efficacy and safety of polyacrylamide hydrogel for the treatment of female stress incontinence: a randomized, prospective, multicenter North American study. *J Urol.* Sep 2014; 192(3): 843-9. PMID 24704117
12. Itkonen Freitas AM, Mentula M, Rahkola-Soisalo P, et al. Tension-Free Vaginal Tape Surgery versus Polyacrylamide Hydrogel Injection for Primary Stress Urinary Incontinence: A Randomized Clinical Trial. *J Urol.* Feb 2020; 203(2): 372-378. PMID 31479396
13. Pai A, Al-Singary W. Durability, safety and efficacy of polyacrylamide hydrogel (Bulkamid((R))) in the management of stress and mixed urinary incontinence: three year follow up outcomes. *Cent European J Urol.* 2015; 68(4): 428-33. PMID 26855795
14. Lose G, Sorensen HC, Axelsen SM, et al. An open multicenter study of polyacrylamide hydrogel (Bulkamid(R)) for female stress and mixed urinary incontinence. *Int Urogynecol J.* Dec 2010; 21(12): 1471-7. PMID 20645077
15. Leone Roberti Maggiore U, Alessandri F, Medica M, et al. Outpatient periurethral injections of polyacrylamide hydrogel for the treatment of female stress urinary incontinence: effectiveness and safety. *Arch Gynecol Obstet.* Jul 2013; 288(1): 131-7. PMID 23371485
16. Mouritsen L, Lose G, Moller-Bek K. Long-term follow-up after urethral injection with polyacrylamide hydrogel for female stress incontinence. *Acta Obstet Gynecol Scand.* Feb 2014; 93(2): 209-12. PMID 24372312
17. Ghoniem G, Corcos J, Comiter C, et al. Cross-linked polydimethylsiloxane injection for female stress urinary incontinence: results of a multicenter, randomized, controlled, single-blind study. *J Urol.* Jan 2009; 181(1): 204-10. PMID 19013613
18. Ghoniem G, Corcos J, Comiter C, et al. Durability of urethral bulking agent injection for female stress urinary incontinence: 2-year multicenter study results. *J Urol.* Apr 2010; 183(4): 1444-9. PMID 20171691
19. Lightner D, Rovner E, Corcos J, et al. Randomized controlled multisite trial of injected bulking agents for women with intrinsic sphincter deficiency: mid-urethral injection of Zuidex via the Implacer versus proximal urethral injection of Contigen cystoscopically. *Urology.* Oct 2009; 74(4): 771-5. PMID 19660800
20. Chapple CR, Haab F, Cervigni M, et al. An open, multicentre study of NASHA/Dx Gel (Zuidex) for the treatment of stress urinary incontinence. *Eur Urol.* Sep 2005; 48(3): 488-94. PMID 15967568
21. Lone F, Sultan AH, Thakar R. Long-term outcome of transurethral injection of hyaluronic acid/dextranomer (NASHA/Dx gel) for the treatment of stress urinary incontinence (SUI). *Int Urogynecol J.* Nov 2010; 21(11): 1359-64. PMID 20571764

MEDICAL POLICY

POLICY TITLE	INJECTABLE BULKING AGENTS FOR THE TREATMENT OF URINARY AND FECAL INCONTINENCE
POLICY NUMBER	MP 4.012

22. Lee PE, Kung RC, Drutz HP. Periurethral autologous fat injection as treatment for female stress urinary incontinence: a randomized double-blind controlled trial. *J Urol.* Jan 2001; 165(1): 153-8. PMID 11125386
23. Bent AE, Tutrone RT, McLennan MT, et al. Treatment of intrinsic sphincter deficiency using autologous ear chondrocytes as a bulking agent. *Neurourol Urodyn.* 2001; 20(2): 157-65. PMID 11170190
24. Strasser H, Marksteiner R, Margreiter E, et al. Autologous myoblasts and fibroblasts versus collagen for treatment of stress urinary incontinence in women: a randomised controlled trial. *Lancet.* Jun 30 2007; 369(9580): 2179-2186. PMID 17604800
25. Kleinert S, Horton R. Retraction--autologous myoblasts and fibroblasts versus collagen [corrected] for treatment of stress urinary incontinence in women: a [corrected] randomised controlled trial. *Lancet.* Sep 06 2008; 372(9641): 789-90. PMID 18774408
26. Peters KM, Dmochowski RR, Carr LK, et al. Autologous muscle derived cells for treatment of stress urinary incontinence in women. *J Urol.* Aug 2014; 192(2): 469-76. PMID 24582537
27. Forte ML, Andrade KE, Butler M, et al. *Treatments for Fecal Incontinence (Comparative Effectiveness Review No. 165).* Rockville, MD: Agency for Healthcare Research and Quality; 2016.
28. Maeda Y, Laurberg S, Norton C. Perianal injectable bulking agents as treatment for faecal incontinence in adults. *Cochrane Database Syst Rev.* Feb 28 2013; (2): CD007959. PMID 23450581
29. Hussain ZI, Lim M, Stojkovic SG. Systematic review of perianal implants in the treatment of faecal incontinence. *Br J Surg.* Nov 2011; 98(11): 1526-36. PMID 21964680
30. Leung FW. Treatment of fecal incontinence - review of observational studies (OS) and randomized controlled trials (RCT) related to injection of bulking agent into peri-anal tissue. *J Interv Gastroenterol.* Oct 2011; 1(4): 202-206. PMID 22586538
31. Graf W, Mellgren A, Matzel KE, et al. Efficacy of dextranomer in stabilised hyaluronic acid for treatment of faecal incontinence: a randomised, sham-controlled trial. *Lancet.* Mar 19 2011; 377(9770): 997-1003. PMID 21420555
32. Dehli T, Stordahl A, Vatten LJ, et al. Sphincter training or anal injections of dextranomer for treatment of anal incontinence: a randomized trial. *Scand J Gastroenterol.* Mar 2013; 48(3): 302-10. PMID 23298304
33. Morris OJ, Smith S, Draganic B. Comparison of bulking agents in the treatment of fecal incontinence: a prospective randomized clinical trial. *Tech Coloproctol.* Oct 2013; 17(5): 517-23. PMID 23525964
34. La Torre F, de la Portilla F. Long-term efficacy of dextranomer in stabilized hyaluronic acid (NASHA/Dx) for treatment of faecal incontinence. *Colorectal Dis.* May 2013; 15(5): 569-74. PMID 23374680
35. Kobashi KC, Albo ME, Dmochowski RR, et al. *Surgical Treatment of Female Stress Urinary Incontinence: AUA/SUFU Guideline.* *J Urol.* Oct 2017; 198(4): 875-883. PMID 28625508
36. National Institute for Health and Care Excellence (NICE). *Urinary incontinence and pelvic organ prolapse in women: management [NG123].* 2019

MEDICAL POLICY

POLICY TITLE	INJECTABLE BULKING AGENTS FOR THE TREATMENT OF URINARY AND FECAL INCONTINENCE
POLICY NUMBER	MP 4.012

37. American College of Obstetricians and Gynecologists (ACOG). Practice Bulletin No. 155: Urinary Incontinence in Women. *Obstet Gynecol.* May 2016;127(5):e66-81. PMID 27548423
38. ACOG Practice Bulletin No. 210: Fecal Incontinence. *Obstet Gynecol.* Apr 2019; 133(4): e260-e273. PMID 30913197
39. Paquette IM, Varma MG, Kaiser AM, et al. The American Society of Colon and Rectal Surgeons' Clinical Practice Guideline for the Treatment of Fecal Incontinence. *Dis Colon Rectum.* Jul 2015; 58(7): 623-36. PMID 26200676
40. Bordeianou, Liliana G. M.D., M.P.H.1; Thorsen, Amy J. M.D.2; Keller, Deborah S. M.S., M.D.3; Hawkins, Alexander T. M.D., M.P.H.4; Messick, Craig M.D.5; Oliveira, Lucia M.D., Ph.D.6; Feingold, Daniel L. M.D.7; Lightner, Amy L. M.D.8; Paquette, Ian M. M.D.9. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Management of Fecal Incontinence. *Diseases of the Colon & Rectum* 66(5):p 647-661, May 2023. | DOI: 10.1097/DCR.0000000000002776
41. National Institute for Health and Care Excellence (NICE). *Injectable bulking agents for faecal incontinence [IPG210].* 2007
42. Centers for Medicare & Medicaid Services. *National Coverage Determination (NCD) for Incontinence Control Devices (230.10).* 1996
43. Blue Cross Blue Shield Association Medical Policy Reference Manual. 7.01.19, *Injectable Bulking Agents for the Treatment of Urinary and Fecal Incontinence.* November 2023.

Other Sources:

1. *Taber's Cyclopedic Medical Dictionary, 19th edition*

X. POLICY HISTORY

[Top](#)

MP 4.012	4/10/20 Consensus review. No change to policy statements. References, rationale and coding updated.
	11/1/21 Minor review. Medically necessary policy statement in men and women with stress urinary incontinence who have failed appropriate conservative therapy expanded to include polyacrylamide hydrogel, which is now FDA approved. Updated: Description/Background and Rationale sections, references and FEP language. Removed dx code N36.41, removed L8603.
	11/23/2022 Consensus review. No change to policy statement. Updated background, references. No coding changes.
	12/18/2023 Consensus review. No change to policy statement. Updated background, references. No coding changes.

[TOP](#)

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company[®], Capital Advantage Assurance Company[®] and Keystone Health Plan[®] Central. Independent licensees of the BlueCross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies