Pediatric COVID-19 Vaccines What Parents/Guardians Should Know

COVID-19 vaccine is recommended for children ages 6 months and up.

Why should children get vaccinated?

Since the beginning of the pandemic, over 15 million children in the United States have gotten COVID-19. While COVID-19 is often milder in children than adults, children can still get very sick and spread it to friends and family.

New COVID-19 variants are currently responsible for most infections and hospitalizations in the United States. Clinical trials proved that this vaccine works very well at preventing serious illness and hospitalizations from COVID-19 in youth.

Vaccination is the best way to keep children healthy and safe.

How do I get my child vaccinated?

Washington state provides all recommended vaccines at no cost for children through age 18. Ask your child's pediatrician or regular clinic if they carry the COVID-19 vaccine.

Families who do not have a health care provider already can call the Help Me Grow WA Hotline at 1-800-322-2588 or go to ParentHelp123.org to find a health care provider, clinic, or other health resources. This service is free and language assistance is available.

You can also visit VaccineLocator.doh.wa.gov and use the filter to see a list of places near you that have the pediatric vaccine.

To get your questions answered: 🗮 Visit: VaccinateWA.org/kids a Call: 1-833-VAX-HELP (1-833-829-4357), Press # Language assistance is available.

How will this help at home?

The COVID-19 vaccine can keep kids protected from the disease, meaning indoor gatherings can be safer to host or attend.

Getting vaccinated Will help children stay safer and protect family members in the household.



Families should continue following other recommended safety measures for the best protection.

Do children need multiple doses?



- Children ages 6 months- 4 years get a 3-dose Pfizer series or a 2-dose Moderna series
- Children ages 5 years and older get a single dose of a Pfizer or Moderna vaccine

Children 6 months+ who are moderately or severely immunocompromised should get a 3-dose series and may get additional doses at least 2 months after their last dose.



Is COVID-19 really a risk to my child?

Although fewer children have been infected with COVID-19 compared to adults, children can:

- Get sick or die from COVID-19
- Spread COVID-19 to others
- Get serious complications from COVID-19, such as "long COVID" or a dangerous inflammatory disease called MIS-C

The vaccine was shown to have very few side effects in this population, and it protects them very well from the disease. The benefits of vaccination far outweigh the risk of getting COVID-19.

Is the vaccine safe for my child?

Yes, the vaccine is considered very safe for children. No serious side effects were detected in clinical trials of the vaccine in youth. The studies are ongoing, and the U.S. has very strong vaccine safety systems to catch any warning signs early.

What are the common side effects of the vaccine in children?

The health risks if children are infected with COVID-19 are much higher than the risk of vaccine side effects.

Like other vaccines, the most common side effects are a sore arm, tiredness, headache, and muscle pain. These symptoms are usually mild.

In <u>clinical trials</u>, children reported side effects were generally mild to moderate in severity and occurred within two days after vaccination, and most went away within one to two days.

What do children need to bring to their appointment?

In most cases, parents and guardians will need to provide consent to vaccination for someone under age 18. They should ask the vaccine location what they need for consent.

I have more questions. Where can I go?

Find answers to more questions at **VaccinateWA.org/kids**. You can also talk to your child's regular health care provider.



DOH 825-013 September 2023

To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email civil.rights@doh.wa.gov